NPIC/TSSG/RED-1896-69 29 October 1969

MEMORANDUM FOR:

Executive Director, NPIC

THROUGH

: Chief, Technical Services & Support Group

Chief, Research & Engineering Division

SUBJECT

: Status Report -- Visual Process Research Program

- The first phase of the Visual Process Research Program is now complete. A wealth of data was collected and analyzed, and many of the findings found application in a variety of situations.
- 2. Primary attention was devoted to assessing the effect of long-term use of microscopes and other optical aids on the visual performance of Center personnel. A previous Center research effort was reviewed by scientists, the results of which indicated that visual acuity scores decreased during a 7 to 14 month time span. The validity of these findings was questioned on several grounds, and the research program was modified in order to:
 - a. test each Center employee who was required to utilize magnification aids in the performance of his duties;
 - b. develop and then follow appropriate test procedures administered by highly skilled test administrators;
 - control for the effects of age on visual performance;
 - d. adhere to a rigid test-retest cycle of 6 months;
 - test for potential "season effects" on visual sensitivity;
 - develop and implement uniform reporting procedures;
 - g. conduct an intensive analysis of the results;

DECLASS REVIEW by NGA/DOD

Excluded from eat: downgrading and 25X1

Approved For Release 2004/02/11 : CIA-RDP78B05703A000100070023-3

SUBJECT: Status Report -- Visual Process Research Program

- h. coordinate Center efforts with the Office of Medical Services and the Center's consulting ophthalmologists;
- i. report the findings to appropriate Center personnel in a timely, understandable fashion.

3. Description of Tests:

- a. Visual acuity, or resolving power,* (monocular and binocular) at near vision (14") and for distance (20')
- b. Vertical phoria* at far (20")
- c. Lateral phoria* at near (14") and at far (20').
- d. Stereo depth* at far (20')
- e. Color discrimination (red-green sensitivity)
- f. Interpupillary distance
- g. Other information:
 - 1.) date of most recent visit to Center's ophthalmologist.
 - 2.) use of eyeglasses
 - 3.) age
 - 4.) job description
 - 5.) general comments

4. Procedures:

Three hundred and sixty four Center employees were divided into six experimental groups, each group representative of the subject population in terms of age (a critical factor in vision research, and one that must be controlled in order to isolate the effects of other variables which might influence visual sensitivity).

^{*} see appendix

Approved For Release 2004/02/11 : CIA-RDP78B05703A000100070023-3

SUBJECT: Status Report -- Visual Process Research Program

Subjects in Group 1 were tested in June, 1968, and retested 6 months later (Dec. 1968). Group 2 subjects were tested in July 1968 and retested in January, 1969, and so forth. All tests were conducted by personnel from the Office of Medical Services/Psychological Services Staff who had been trained by the author. The Office of Medical Services nurse assigned to the Center was responsible for all scheduling & record keeping.

- 5. Results: The several analyses of the results indicated:
 - a. no significant differences in scores on any parameter among the 6 groups during the initial 6 month test phase;
 - b. no significant differences in scores on any parameter among the same 6 groups during the 6 month retest phase;
 - no significant differences in test vs. retest scores on any parameter for the results of each group analyzed independently;
 - d. group averages were at least comparable, if not superior, to national norms for all tests except lateral phoria at 20', where a statistically significant exophoria* of 2.0 to 2.5 prism diopters was found. No major significance was placed in this finding by the several experts consulted. Furthermore, the results agree with data supplied to the Center by U.S. Naval School of Aviation Medicine.
 - e. average interpupillary distances was approximately 64 mm., but the range of scores indicated that a few subjects would find it difficult if not impossible to use binocular instruments;
 - f. fewer than 40 subjects were found to be red-green color deficient;
 - g. the vast majority of the subjects were capable of resolving the most difficult stereo test in the test instrument;

3.2

Approved or Release 2004/02/11 : CIA-RDP78B 2003A000100070023-3

SUBJECT: Status Report -- Visual Process Research Program

- h. no effects of season on visual acuity were apparent, unlike the situation where the amount of sunlight associated with seasons was found by the Navy to influence visual sensitivity (dark adaptation);
- i. of all the interpreters who had prescription eyeglasses (some 65% of the subject population), approximately 75% of them wore their glasses when using magnification aids to interpret imagery (a matter of convenience), despite the fact that the microscopes can correct for refractive errors, but not astigmatism.

6. Selected Application of Results:

- a. Earlier indications that continued use of microscopes was having a deleterious effect on visual acuity were not corroborated in the present study, suggesting that inappropriate test procedures were followed in the previous test program.
- b. The names and records of the 40 individuals who were judged by the author to demonstrate the poorest visual performance in the test program were submitted to the Center's consultant ophthalmologist for further review. Results are expected within 30 days, according to an Office of Medical Services spokesman.
- c. Once that information is received, work will be devoted to further specifying the visual requirements for photo interpretation. The Office of Medical Services has agreed to provide detailed vision information to the Center on all applicants.
- d. Visual acuity scores were correlated with mensuration accuracy. _____ included this analysis in their mensuration research program, and the results indicated that the two measures are not related -- due primarily to the fact that the photogrammetrist are typically well corrected and can compensate for any residual refractive errors with adjustments available on their comparator optics.
- e. The IEG was recently provided the names of 38 interpreters who were judged by the author to be at least marginally deficient in color perception. Very liberal criteria were employed in selecting these people, and the fact that one's

SUBJECT: Status Report -- Visual Process Research Program

25X

25X

name appears on the list need not necessarily indicate a true color deficiency. The information was provided to insure that IEG management is aware of color deficient PIs who might be tasked with the exploitation of color materials, or worse, with the determination of the intelligence value of color imagery. Stimulus materials for a more sensitive color test have been secured, and a follow up color test will be administered if so requested by IEG.

1 lits

f. A rather extensive review of the literature on color vision has been in progress by the author for some time. This effort is directed toward the exploitation of color materials, and is in support of one portion of the color-oriented research program in RED under the direction of

25X1

g. A stereo test more sensitive than the one employed in the survey was developed by for the Microscope Design Criteria study. The test was recently evaluated by this office as a means of selecting photogrammetrists with particularly high stereo acuity. The results of the preliminary test are encouraging, and the recommendation has been made to continue employing the test as a screening device. The stimulus materials utilized in the tests are presently being calibrated by APSD.

25X1

h. One early product of the vision survey was an age/job description profile for the Center personnel who participated in the program. The information is now of historical use only, and it should be updated to reflect the organizational & personnel changes that have occurred in the past year or so.

ok-doe

i. Several commercial firms that employ large numbers of microscopists were asked if long term use of magnification aids influenced visual sensitivity. A surprisingly detailed and candid reply was received from which indicated that "...no noticable eye deterioration exists as the result of long use of microscope. On the contrary, doctors are convinced this tends to stall normal age-related eye deterioration."

25X1

j. The plight of several IEG interpreters with unusually small interpupillary distances was discussed with IEG management,

Approve pr Release 2004/02/11 : CIA-RDP78B 703A000100070023-3

SUBJECT: Status Report -- Visual Process Research Program

since it was determined that it would be difficult if not virtually impossible for them to utilize binocular instruments due to a minimum instrument IPD of approximately 55 mm. Similar information was provided RED personnel concerned with microscope design.

h. Results from the survey concerned with the use of eyeglasses when viewing through microscopes were provided to RED personnel tasked with designing microscopes which have, by virtue of elaborate optical characteristics, an exceedingly small exit pupil. The exit pupil in one system is so limited that it would be impossible for an interpreter to wear corrections other than contact lenses. On the basis of the survey results, it was concluded that acuity adapters should be provided for at least those interpreters afflicted with marked astigmatism.

oldes Loss.

7. HFS/ATB/RED is	currently	drafting	a detailed	_report	
on the Visual Process Research Program.				1	
				•	
•					

Distribution:

25X1

Orig - Addressee

- 1 TSSG
- 1 TSSG/RED
- 2 RED/ATB/HFS

.

Chief, Human Factors Section, ATB

25X

25

TSSG/ESD-1/70

MEMORANDUM FOR: Executive Director, NPIC

SUBJECT

: Response to Inquiry Regarding Illumination and

Temperature Stability of Light Tables

1. The summary section of T&E Report No. 69-06 on the prototype _____1540 Light Table states: "The light table takes approximately four hours to achieve illumination and temperature stability." Although the statement as it stands can be taken as an implication that the warm-up time is excessive, the light table is completely usable long before stability is reached. The warm-up characteristic curves on pages 24 and 25 show that the illumination was well over 50% of full intensity when first turned on, reached approximately 90% of full intensity within 15 minutes and, for all practical purposes, reached full intensity within 30 minutes.

- 2. The light intensity stabilized completely after two hours, but the film stage temperature continued to rise slowly before stabilizing after four hours. Stabilized illumination and temperature values were measured for comparison with design objectives as stated on pages 10 and 11 of the report.
- 3. Three other prototype light tables were tested by TEB/ESD this past year. They were all air-cooled and required from 15 minutes to 30 minutes to reach full light intensity and 3 to 6 hours to reach temperature stability. Therefore, the liquid-cooled table is not at all unusual in its performance regarding these factors.

Chief, Technical Services & Support Group NPIC

Distribution:

БХ1

Original - Addressee

2 - NPIC/TSSG

1 - NPIC/TSSG/ESD

roved For Release 2004/02/1 MglA-7073B05703A0) 0010
TRANSMITTAL SLIP	
TO:	
ROOM NO.	
REMARKS:	1
Pla Drepare a response	
Jar	
	- I
to see the response.	
to see the response.	
FROM: EXEL TSSQ	
ROOM NO. BUILDING EXTENSION	1
rov <mark>ed For Releas</mark> e 2004/02/11 : CIA-RDP78B05703A0	<u>)</u> 010
FORM NO . 241 REPLACES FORM 36-8 WHICH MAY BE USED.	7)

	Marie Colombia Colomb	о	
		23 . Mac. &	of Control of the Con
1			
	lave read	the THE A	Property of
1 con the		1540 light	f.
		ic to be abs	To the second
	e liekt tood		
Meanin	atoly from	Lours to 201	
Monda.	2.1.1).	perefere sta	Solo Por
	4217	esperiod us	real
for other	licht ta	bles?	,
			2

Approved For Release 2004/02/11 : 6/A-RDP78B05703A00010b070023

LIKE TO BE BRIFTED

BY ON THE

UISUAL PROCESS RESHARCH

PROGRAM

2. JTH ASKUD TWO
QUESTIONS ON THE 29
OCT UPP PROGRAM
SMATUS PUPOPT

JOB DISCR. PROFILE?

も、いけい PDOUIDES ACUITY
Approper 1705 1705 2904 102/11: CIA-RDP78B05703A000100070023-1